


## pTagFP635-Cx26 vector

The vector sequence has been compiled using the information from sequence databases, published literature, and other sources, together with partial sequences obtained by Evrogen. This vector has not been completely sequenced.



For vector sequence, please visit our Web site at <http://www.evrogen.com/support/vector-info.shtml>

### Location of features

PCMV IE: 1-589  
 Enhancer region: 59-465  
 TATA box: 554-560  
 Transcription start point: 583  
 Connexin 26: 683-1360  
 TagFP635: 1382-2095  
 SV40 early mRNA polyadenylation signal  
 Polyadenylation signals: 2248-2253 & 2277-2282  
 mRNA 3' ends: 2286 & 2298  
 f1 single-strand DNA origin: 2345-2800  
 Bacterial promoter for expression of Kan<sup>r</sup> gene  
 -35 region: 2862-2867; -10 region: 2885-2890  
 Transcription start point: 2897  
 SV40 origin of replication: 3141-3276  
 SV40 early promoter  
 Enhancer (72-bp tandem repeats): 2974-3045 & 3046-3117  
 21-bp repeats: 3121-3141, 3142-3162 & 3164-3184  
 Early promoter element: 3197-3203  
 Major transcription start points: 3193, 3231, 3237 & 3242  
 Kanamycin/neomycin resistance gene  
 Neomycin phosphotransferase coding sequences:  
 Start codon (ATG): 3325-3327; Stop codon: 4117-4119  
 G->A mutation to remove Pst I site: 3507  
 C->A (Arg to Ser) mutation to remove BssH II site: 3853  
 Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal  
 Polyadenylation signals: 4355-4360 & 4368-4373  
 pUC plasmid replication origin: 4704-5347

Product	Cat.#	Size
pTagFP635-Cx26 vector	<b>FP382</b>	20 µg
The price does not include delivery. The price varies in different countries. Please contact your local distributor for exact prices and delivery information.		
Vector type	mammalian expression vector	
Reporter	TagFP635	
Reporter codon usage	mammalian	
Promoter for TagFP635	P <sub>CMV IE</sub>	
Host cells	mammalian	
Selection	prokaryotic - kanamycin eukaryotic - neomycin (G418)	
Replication	prokaryotic - pUC ori eukaryotic - SV40 ori	
Use	far-red fluorescent labeling of connexin 26	

### Vector description

pTagFP635-Cx26 is a mammalian expression vector encoding TagFP635-Cx26 fusion protein. The vector can be used for fluorescent labeling of connexin 26 in living cells.

TagFP635 codon usage is optimized for high expression in mammalian cells, i.e. humanized (Haas et al. 1996). Rat connexin 26 is fused to the TagFP635 N-terminus.

pTagFP635-Cx26 can be used as a source of TagFP635-Cx26 hybrid sequence. The vector backbone contains unique restriction sites that permit its excision and further insertion into expression vector of choice.

**Note:** The plasmid DNA was isolated from dam<sup>+</sup>-methylated *E. coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a dam<sup>-</sup> host and make fresh DNA.

The vector backbone also contains immediate early promoter of cytomegalovirus (P<sub>CMV IE</sub>) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli*, and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3' end of the reporter mRNA.

SV40 early promoter (P<sub>SV40</sub>) provides neomycin resistance gene (Neo<sup>r</sup>) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan<sup>r</sup>) in *E. coli*. Kan<sup>r</sup>/Neo<sup>r</sup> gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

### Expression in mammalian cells

pTagFP635-Cx26 can be transfected into mammalian cells by any known transfection method. CMV promoter provides strong, constitutive expression of the TagFP635-Cx26 fusion in eukaryotic cells. If required, stable transformants can be selected using G418 [Gorman 1985].

### Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.

### References

- Gorman (1985). "High efficiency gene transfer into mammalian cells." In: *DNA cloning: A Practical Approach, Vol. II*. Ed. by Glover. (IRL Press, Oxford, U.K.) Pp. 143-190.
- Haas et al. (1996) "Codon usage limitation in the expression of HIV-1 envelope glycoprotein." *Curr Biol*, 6 (3): 315-324 / pmid: 8805248

### Notice to Purchaser:

Evrogen Fluorescent Protein Products (the Products) are intended for research use only. The Products are covered by U.S. Pat. 7,417,131 and other Evrogen Patents and/or Patent applications pending. By use of these Products, you accept the terms and conditions of the applicable Limited Use Label License.

The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

**MATERIAL SAFETY DATA SHEET INFORMATION:** To the best of our knowledge, these products do not require a Material Safety Data Sheet. However, all the properties of these products (and, if applicable, each of their components) have not been thoroughly investigated. Therefore, we recommend that you use gloves and eye protection, and wear a laboratory coat when working with these products.