


## pTagFP635-laminB1 vector

The vector sequence has been compiled using the information from sequence databases, published literature, and other sources, together with partial sequences obtained by Evrogen. This vector has not been completely sequenced.



For vector sequence, please visit our Web site at <http://www.evrogen.com/support/vector-info.shtml>

### Location of features

**P<sub>CMV IE</sub>:** 1-589  
**Enhancer region:** 59-465  
**TATA box:** 554-560  
**Transcription start point:** 583  
**Kozak consensus translation initiation site:** 606-616  
**TagFP635**  
**Start codon (ATG):** 613-615  
**Last amino acid in TagFP635:** 1321-1323  
**Lamin B1:** 1354-3114  
**Stop codon:** 3112-3114  
**SV40 early mRNA polyadenylation signal**  
**Polyadenylation signals:** 3275-3280 & 3304-3309  
**mRNA 3' ends:** 3313 & 3325  
**f1 single-strand DNA origin:** 3372-3827  
**Bacterial promoter for expression of Kan<sup>r</sup> gene**  
**-35 region:** 3889-3894; **-10 region:** 3912-3917  
**Transcription start point:** 3924  
**SV40 origin of replication:** 4168-4303  
**SV40 early promoter**  
**Enhancer (72-bp tandem repeats):** 4001-4072 & 4073-4144  
**21-bp repeats:** 4148-4168, 4169-4189 & 4191-4211  
**Early promoter element:** 4224-4230  
**Major transcription start points:** 4220, 4258, 4264 & 4269  
**Kanamycin/neomycin resistance gene**  
**Neomycin phosphotransferase coding sequences:**  
**Start codon (ATG):** 4352-4354; **Stop codon:** 5144-5146  
**G->A mutation to remove Pst I site:** 4534  
**C->A (Arg to Ser) mutation to remove BssH II site:** 4880  
**Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal**  
**Polyadenylation signals:** 5382-5387 & 5395-5400  
**pUC plasmid replication origin:** 5731-6374

Product	Cat.#	Size
pTagFP635-laminB1 vector	<b>FP387</b>	20 µg

The price does not include delivery. The price varies in different countries. Please contact your local distributor for exact prices and delivery information.

Vector type	mammalian expression vector
Reporter	TagFP635
Reporter codon usage	mammalian
Promoter for TagFP635	P <sub>CMV IE</sub>
Host cells	mammalian
Selection	prokaryotic - kanamycin eukaryotic - neomycin (G418)
Replication	prokaryotic - pUC ori eukaryotic - SV40 ori
Use	far-red fluorescent labeling of lamin B1

### Vector description

pTagFP635-laminB1 is a mammalian expression vector encoding TagFP635-lamin B1 fusion protein. The vector can be used for fluorescent labeling of lamin B1 in living cells.

TagFP635 codon usage is optimized for high expression in mammalian cells, i.e. humanized [Haas et al. 1996]. Human lamin B1 is fused to the TagFP635 C-terminus. To increase mRNA translation efficiency, Kozak consensus translation initiation site is generated upstream of TagFP635-lamin B1 coding sequence [Kozak 1987].

pTagFP635-laminB1 can be used as a source of TagFP635-lamin B1 hybrid sequence. The vector backbone contains unique restriction sites that permit it excision and further insertion into expression vector of choice.

**Note:** The plasmid DNA was isolated from dam<sup>+</sup>-methylated *E. coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a dam<sup>-</sup> host and make fresh DNA.

The vector backbone also contains immediate early promoter of cytomegalovirus (P<sub>CMV IE</sub>) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli*, and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3' end of the reporter mRNA.

SV40 early promoter (P<sub>SV40</sub>) provides neomycin resistance gene (Neo<sup>r</sup>) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan<sup>r</sup>) in *E. coli*. Kan<sup>r</sup>/Neo<sup>r</sup> gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

### Expression in mammalian cells

pTagFP635-laminB1 can be transfected into mammalian cells by any known transfection method. CMV promoter provides strong, constitutive expression of the TagFP635-lamin B1 fusion in eukaryotic cells. If required, stable transformants can be selected using G418 [Gorman 1985].

### Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.

### References

- Gorman (1985). "High efficiency gene transfer into mammalian cells." In: *DNA cloning: A Practical Approach, Vol. II*. Ed. by Glover. (IRL Press, Oxford, U.K.) Pp. 143-90.
- Haas et al. (1996) "Codon usage limitation in the expression of HIV-1 envelope glycoprotein." *Curr Biol*, 6 (3): 315-24 / pmid: 8805248
- Kozak (1987) "An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs." *Nucleic Acids Res*, 15 (20): 8125-48 / pmid: 3313277

### Notice to Purchaser:

Evrogen Fluorescent Protein Products (the Products) are intended for research use only and covered by Evrogen Patents and/or Patent applications pending. By use of these products, you accept the terms and conditions of the applicable Limited Use Label License (enclosed).

The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

**MATERIAL SAFETY DATA SHEET INFORMATION:** To the best of our knowledge, these products do not require a Material Safety Data Sheet. However, all the properties of these products (and, if applicable, each of their components) have not been thoroughly investigated. Therefore, we recommend that you use gloves and eye protection, and wear a laboratory coat when working with these products.