



## Anti-KillerRed antibody

Product	Cat.#	Lot.#	Size
Anti-KillerRed antibody	<b>AB961</b>	96101240513	100 µg

### Use

- Western blot
- Immunoblotting
- ICC
- ELISA



Western blot detection of fluorescent proteins using anti-KillerRed antibody.

### Description

Rabbit polyclonal antibody against KillerRed, KillerOrange, ArrestRed, and JRed.

**Specificity:** The antibody has been selected to recognize both denatured and native KillerRed, KillerOrange, JRed and ArrestRed. The antibody shows little or no cross-reactivity with other fluorescent proteins tested.

**Immunogen:** Full-length recombinant denatured KillerRed.

**Antibody preparation:** Full-length recombinant KillerRed was purified from transformed *E. coli* using organic extraction and hydrophobic chromatography. Antibody was produced in rabbits immunized with the recombinant denatured KillerRed and purified by KillerRed affinity chromatography.

**Formulation:** Lyophilized from the PBS buffer containing 0.05% NaN<sub>3</sub> and 0.5% trehalose; pH 7.4.

**Reconstitution:** Reconstitute with sterile water or 50% glycerol to a concentration of 1 mg/ml.

**Storage:** Lyophilized samples are stable for twelve months from date of receipt when stored at -20 °C. The presence of silica gel drier is advisable.

Reconstituted with sterile water, antibody can be stored at 2 - 8 °C three months without detectable loss of activity.

Reconstituted with 50% glycerol, antibody can be stored at -20 °C in a manual defrost freezer for six months without detectable loss of activity. Aliquot antibody upon reconstitution. Avoid repeated freeze / thaw cycles.

### Recommendations for use

The antibody can be used to recognize KillerRed, KillerOrange, JRed, ArrestRed and their fusions.

#### Working concentrations:

For Western blot use at a dilution of 1 : 1 0000;

For ELISA use at a dilution of 1 : 20 000;

For immunocytochemistry use at a dilution of 1 : 5 000.

**Note.** Optimal dilutions/concentrations should be determined by the end user.

**Tissue (cells) fixation for immunohistochemistry:** Formaldehyde (formalin, paraform) fixation is recommended. For example, tissues can be fixed in PBS containing 4% formaldehyde for 10-15 min, treated with 0.1% saponin in PBS for 10-15 min, and washed three times in PBS.

**Sample preparation for Western blot:** To a sample containing 10-100 ng of a target protein, add an equal volume of 2X SDS-PAGE sample buffer. Heat the sample at 95 °C before loading on a gel or spotting on a membrane (for dots).

### Notice to Purchaser:

These products are intended for research purposes only.

**MSDS information** is available at <http://evrogen.com/support/MSDS-info.shtml>