



## pTagGFP2-C vector

The vector sequence has been compiled using the information from sequence databases, published literature, and other sources, together with partial sequences obtained by Evrogen. This vector has not been completely sequenced.



For vector sequence, please visit our Web site at <http://www.evrogen.com/products/vectors.shtml>

### Multiple cloning site (MCS)



\* — not unique sites.

# — sites are blocked by *dam* methylation. If you wish to digest the vector with these enzymes, you will need to transform the vector into a *dam*<sup>-</sup> host and make fresh DNA.

### Location of features

P<sub>CMV IE</sub>: 1-589  
 Enhancer region: 59-465  
 TATA box: 554-560  
 Transcription start point: 583  
 TagGFP2  
 Kozak consensus translation initiation site: 600-610  
 Start codon (ATG): 607-609; Stop codon: 1399-1401  
 Last amino acid in TagGFP2: 1318-1320  
 MCS: 1321-1398  
 SV40 early mRNA polyadenylation signal  
 Polyadenylation signals: 1541-1546 & 1570-1575  
 mRNA 3' ends: 1579 & 1591  
 f1 single-strand DNA origin: 1638-2093  
 Bacterial promoter for expression of Kan<sup>r</sup> gene  
 -35 region: 2155-2160; -10 region: 2178-2183  
 Transcription start point: 2190  
 SV40 origin of replication: 2434-2569  
 SV40 early promoter  
 Enhancer (72-bp tandem repeats): 2267-2338 & 2339-2410  
 21-bp repeats: 2414-2434, 2435-2455 & 2457-2477  
 Early promoter element: 2490-2496  
 Major transcription start points: 2486, 2524, 2530 & 2535  
 Kanamycin/neomycin resistance gene  
 Neomycin phosphotransferase coding sequences:  
 Start codon (ATG): 2618-2620; Stop codon: 3410-3412  
 G->A mutation to remove Pst I site: 2800  
 C->A (Arg to Ser) mutation to remove BssH II site: 3146  
 Herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signal  
 Polyadenylation signals: 3648-3653 & 3661-3666  
 pUC plasmid replication origin: 3997-4640

### References

- Gorman, C. (1985). "High efficiency gene transfer into mammalian cells." In: *DNA cloning: A Practical Approach, Vol. II*. Ed. by Glover. (IRL Press, Oxford, U.K.) Pp. 143-190.
- Haas, J. et al. (1996) "Codon usage limitation in the expression of HIV-1 envelope glycoprotein." *Curr Biol*, 6 (3): 315-324 / pmid: 8805248
- Kozak, M. (1987) "An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs." *Nucleic Acids Res*, 15 (20): 8125-8148 / pmid: 3313277

### Notice to Purchaser:

TagGFP2-related materials (also referred to as "Products") are intended for research use only. The Products are covered by U.S. Pat. 7,417,131; 7,605,230; 7,888,113; European Pat. 06809023; and other Evrogen Patents and/or Patent applications pending. By use of these Products, you accept the terms and conditions of the applicable Limited Use Label License #001: <http://www.evrogen.com/products/Evrogen-FP-license.shtml>. The CMV promoter is covered under U.S. Patents 5,168,062 and 5,385,839, and its use is permitted for research purposes only. Any other use of the CMV promoter requires a license from the University of Iowa Research Foundation, 214 Technology Innovation Center, Iowa City, IA 52242.

**MSDS information** is available at <http://www.evrogen.com/MSDS.shtml>

Product	Cat.#	Size
pTagGFP2-C vector	<b>FP191</b>	20 µg
Vector type	mammalian expression vector	
Reporter	TagGFP2	
Reporter codon usage	mammalian	
Promoter for TagGFP2	P <sub>CMV IE</sub>	
Host cells	mammalian	
Selection	prokaryotic - kanamycin eukaryotic - neomycin (G418)	
Replication	prokaryotic - pUC ori eukaryotic - SV40 ori	
Use	TagGFP2 expression in mammalian cells; generation of fusions to the TagGFP2 C-terminus	

### Vector description

pTagGFP2-C is a mammalian expression vector encoding green fluorescent protein TagGFP2. The vector allows generation of fusions to the TagGFP2 C-terminus and expression of TagGFP2 fusions or TagGFP2 alone in eukaryotic (mammalian) cells.

TagGFP2 codon usage is optimized for high expression in mammalian cells (humanized) [Haas et al. 1996]. To increase mRNA translation efficiency, Kozak consensus translation initiation site is generated upstream of the TagGFP2 coding sequence [Kozak 1987]. Multiple cloning site (MCS) is located between TagGFP2 coding sequence and SV40 polyadenylation signal (SV40 polyA).

The vector backbone contains immediate early promoter of cytomegalovirus (P<sub>CMV IE</sub>) for protein expression, SV40 origin for replication in mammalian cells expressing SV40 T-antigen, pUC origin of replication for propagation in *E. coli*, and f1 origin for single-stranded DNA production. SV40 polyadenylation signals (SV40 poly A) direct proper processing of the 3'-end of the reporter mRNA.

SV40 early promoter (P<sub>SV40</sub>) provides neomycin resistance gene (Neo<sup>r</sup>) expression to select stably transfected eukaryotic cells using G418. Bacterial promoter (P) provides kanamycin resistance gene expression (Kan<sup>r</sup>) in *E. coli*. Kan<sup>r</sup>/Neo<sup>r</sup> gene is linked with herpes simplex virus (HSV) thymidine kinase (TK) polyadenylation signals.

### Generation of TagGFP2 fusion proteins

A localization signal or a gene of interest can be cloned into MCS of the vector. It will be expressed as a fusion to the TagGFP2 C-terminus when inserted in the same reading frame as TagGFP2 and no in-frame stop codons are present. TagGFP2-tagged fusions retain fluorescent properties of the native protein allowing fusion localization *in vivo*. Unmodified vector will express TagGFP2 when transfected into eukaryotic (mammalian) cells.

**Note:** The plasmid DNA was isolated from *dam*<sup>+</sup>-methylated *E. coli*. Therefore some restriction sites are blocked by methylation. If you wish to digest the vector using such sites you will need to transform the vector into a *dam*<sup>-</sup> host and make fresh DNA.

### Expression in mammalian cells

pTagGFP2-C vector can be transfected into mammalian cells by any known transfection method. CMV promoter provides strong, constitutive expression of TagGFP2 or its fusions in eukaryotic cells. If required, stable transformants can be selected using G418 [Gorman 1985].

### Propagation in *E. coli*

Suitable host strains for propagation in *E. coli* include DH5alpha, HB101, XL1-Blue, and other general purpose strains. Plasmid incompatibility group is pMB1/ColE1. The vector confers resistance to kanamycin (30 µg/ml) to *E. coli* hosts. Copy number in *E. coli* is about 500.